

AMPA Receptor 4 (GluA 4) (2F9) Rabbit Monoclonal Antibody

AMPA Receptor 4 (GluA 4) (2F9) Rabbit Monoclonal Antibody
Catalog # AP93742

Product Information

Application	WB, FC
Primary Accession	P48058 , Q9Z2W8 , P19493
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Calculated MW	100871

Additional Information

Gene ID	2893
Dilution	WB~~1:1000 FC~~1:10~50
Storage Conditions	-20°C


Protein Information

Name	GRIA4 {ECO:0000303 PubMed:29220673, ECO:0000312 HGNC:HGNC:4574}
Function	Ionotropic glutamate receptor that functions as a ligand- gated cation channel, gated by L-glutamate and glutamatergic agonists such as alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), quisqualic acid, and kainic acid (By similarity). L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system and plays an important role in fast excitatory synaptic transmission (By similarity). Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse upon entry of monovalent and divalent cations such as sodium and calcium. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist (By similarity). In the presence of CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of L-glutamate (PubMed: 21172611).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P19493}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19493} Postsynaptic cell membrane {ECO:0000250 UniProtKB:P19493}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19493}. Cell projection, dendrite {ECO:0000250 UniProtKB:P19493}. Postsynaptic cell membrane {ECO:0000250 UniProtKB:P42262}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P42262}

Background

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes composed of multiple subunits, arranged to form ligand-gated ion channels. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. The subunit encoded by this gene belongs to a family of AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate)-sensitive glutamate receptors, and is subject to RNA editing (AGA->GGA; R->G). Alternative splicing of this gene results in transcript variants encoding different isoforms, which may vary in their signal transduction properties. Some haplotypes of this gene show a positive association with schizophrenia. [provided by RefSeq, Jul 2008]

Images



Western blot analysis of extracts from Rat brain tissue using AP93742 at 1:1000.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.