

CCL26 Rabbit pAb

CCL26 Rabbit pAb
Catalog # AP93948

Product Information

Application	IHC-P, IHC-F, IF
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from rat CCL26
Epitope Specificity	21-94/94
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene is one of two Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 7. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for normal peripheral blood eosinophils and basophils. The product of this gene is one of three related chemokines that specifically activate chemokine receptor CCR3. This chemokine may contribute to the eosinophil accumulation in atopic diseases. [provided by RefSeq, Jul 2008]

Additional Information


Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1ug/test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images

Blank control:U-2OS.



Primary Antibody (green line): Rabbit Anti-CCL26 antibody (AP93948)

Dilution: 1 μ g /10⁶ cells;


Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE


Dilution: 1 μ g /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 20% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Paraformaldehyde-fixed, paraffin embedded (rat thyroid gland tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CCL26) Polyclonal Antibody, Unconjugated (AP93948) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control:U-2OS. Primary Antibody (green line):

Rabbit Anti-CCL26 antibody (AP93948) Dilution: 1 μ g

/10⁶ cells; Isotype Control Antibody (orange line): Rabbit

IgG . Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 μ g /test. Protocol The cells were fixed with 4%

PFA (10min at room temperature) and then permeabilized

with 20% PBST for 20 min at room temperature. The cells

were then incubated in 5%BSA to block non-specific

protein-protein interactions for 30 min at room

temperature. Cells stained with Primary Antibody for 30

min at room temperature. The secondary antibody used

for 40 min at room temperature. Acquisition of 20,000

events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.