

Neuronal thread protein AD7c-NTP Rabbit pAb

Neuronal thread protein AD7c-NTP Rabbit pAb

Catalog # AP93969

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Neuronal thread protein AD7c-NTP
Epitope Specificity	301-375/375
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	AD7c-NTP is detected in increased concentration in the cortical neurons, brain-tissue extracts, cerebrospinal fluid, and urine early in the course of AD neurodegeneration, and its level is positively correlated with the severity of dementia. All these characteristics make it a possible biomarker for AD.

Additional Information


Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1ug/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images


Blank control: SHSY5Y.
Primary Antibody (green line): Rabbit Anti-Neuronal thread protein AD7c-NTP antibody (AP93969)
Dilution: 1ug/Test;
Secondary Antibody : Goat anti-rabbit IgG-FITC



Dilution: 0.5ug/Test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.




Sample: Brain (Mouse) Lysate at 40 ug

Primary: Anti- AD7c-NTP(AP93969) at 1/300 dilution

Secondary: HRP conjugated Goat-Anti-rabbit IgG (AP93969-HRP) at 1/5000 dilution

Predicted band size: 41 kD

Observed band size: 41 kD



Tissue/cell: human brain tissue; 4%

Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min;

Incubation: Anti-AD7c-NTP Polyclonal Antibody, Unconjugated(AP93969) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.