

HPV16 E6 + HPV18 E6 Rabbit pAb

HPV16 E6 + HPV18 E6 Rabbit pAb

Catalog # AP93982

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	HPV18, HPV16
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from HPV16 E6+ HPV18 E6
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


Additional Information

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1ug /Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background


This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Blank control(black line):Hela.
Primary Antibody (green line): Rabbit Anti-HPV16 E6 + HPV18 E6 antibody (AP93982)
Dilution:1ug/Test;
Secondary Antibody(white blue line): Goat anti-rabbit IgG-AF488
Dilution: 0.5ug/Test.
Isotype control(orange line): Normal Rabbit IgG
Protocol
The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then

incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Sample:

Lane 1: Recombinant HPV16 E6 protein, DsbC &

His(AP93982) 0.1ug

Lane 2: Recombinant HPV16 E6 protein, DsbC &


His(AP93982) 0.05ug

Primary: Anti-HPV16 E6 + HPV18 E6 (AP93982) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 11 kDa

Observed band size: 50 kDa



Tissue/cell: human colon carcinoma; 4%

Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling

bathing for 15min; Block endogenous peroxidase by 3%

Hydrogen peroxide for 30min; Blocking buffer (normal

goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-HPV16 E6+HPV18 E6 Polyclonal Antibody,

Unconjugated(AP93982) 1:200, overnight at 4°C, followed

by conjugation to the secondary antibody(SP-0023) and

DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.