

p19 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9406A

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	P55273
Other Accession	Q60773
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	17700
Antigen Region	15-43

Additional Information

Gene ID	1032
Other Names	Cyclin-dependent kinase 4 inhibitor D, p19-INK4d, CDKN2D
Target/Specificity	This p19 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 15-43 amino acids from the N-terminal region of human p19.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	p19 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CDKN2D
Function	Interacts strongly with CDK4 and CDK6 and inhibits them.
Cellular Location	Nucleus. Cytoplasm


Background

p19 is a member of the INK4 family of cyclin-dependent kinase inhibitors. This protein has been shown to form a stable complex with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. The abundance of the transcript of this gene was found to oscillate in a cell-cycle dependent manner with the lowest expression at mid G1 and a maximal expression during S phase. The negative regulation of the cell cycle involved in this protein was shown to participate in repressing neuronal proliferation, as well as spermatogenesis.


References

Cunningham, J.M., et al. Br. J. Cancer 101(8):1461-1468(2009)
Zhang, W., et al. Acta Biochim. Biophys. Sin. (Shanghai) 41(5):414-428(2009)
Agarwal, S.K., et al. J. Clin. Endocrinol. Metab. 94(5):1826-1834(2009)
Goode, E.L., et al. Cancer Epidemiol. Biomarkers Prev. 18(3):935-944(2009)
Mavaddat, N., et al. Cancer Epidemiol. Biomarkers Prev. 18(1):255-259(2009)


Images



Anti-p19 Antibody (N-term) at 1:2000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



p19 Antibody (N-term)(Cat. #AP9406a) IHC analysis in formalin fixed and paraffin embedded testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the p19 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



p19 Antibody (N-term) (Cat. #AP9406a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.