

# AMPK alpha-1 Rabbit pAb

AMPK alpha-1 Rabbit pAb  
Catalog # AP94150

## Product Information

---

<b>Application</b>	WB, E
<b>Reactivity</b>	Human
<b>Predicted</b>	Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	64 kDa
<b>Physical State</b>	Liquid
<b>Immunogen</b>	Recombinant human AMPK alpha-1 protein
<b>Epitope Specificity</b>	1-207/559
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Cytoplasm. Nucleus. Note=In response to stress, recruited by p53/TP53 to specific promoters.
<b>SIMILARITY</b>	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. SNF1 subfamily. Contains 1 protein kinase domain.
<b>SUBUNIT</b>	AMPK is a heterotrimer of an alpha catalytic subunit (PRKAA1 or PRKAA2), a beta (PRKAB1 or PRKAB2) and a gamma non-catalytic subunits (PRKAG1, PRKAG2 or PRKAG3). Interacts with FNIP1 and FNIP2.
<b>Post-translational modifications</b>	Ubiquitinated. Phosphorylated at Thr-183 by STK11/LKB1 in complex with STE20-related adapter-alpha (STRADA) pseudo kinase and CAB39. Also phosphorylated at Thr-183 by CAMKK2; triggered by a rise in intracellular calcium ions, without detectable changes in the AMP/ATP ratio. CAMKK1 can also phosphorylate Thr-183, but at much lower level. Dephosphorylated by protein phosphatase 2A and 2C (PP2A and PP2C). Phosphorylated by ULK1 and ULK2; leading to negatively regulate AMPK activity and suggesting the existence of a regulatory feedback loop between ULK1, ULK2 and AMPK.
<b>DISEASE</b>	Defects in CRYAB are the cause of myofibrillar alpha-B crystallin-related (MFM-CRYAB) [MIM:608810]. A neuromuscular disorder that results in weakness of the proximal and distal limb muscles, weakness of the neck, velopharynx and trunk muscles, hypertrophic cardiomyopathy, and cataract in a subset of patients.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]

## Additional Information


<b>Target/Specificity</b>	Lens as well as other tissues.
<b>Dilution</b>	WB=1:500-2000,ELISA=1:5000-10000
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Background


The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]

## Images

Rabbit Anti-Human AMPK alpha-1 Antibody Bind with Human AMPK alpha-1 protein, His Tag




Measured by its binding ability in a indirect ELISA. Immobilized Human AMPK alpha-1 protein, His Tag (Cat. AP94150) at 2 µg/mL (100 µL/well) can bind Rabbit Anti-Human AMPK alpha-1 Antibody, the EC50 is 133.7 ng/mL.




Sample:  
Lane 1: Human HUVEC cell lysates  
Lane 2: Human SH-SY5Y cell lysates  
Primary: Anti-AMPK alpha-1 (AP94150) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 64 kDa  
Observed band size: 64 kDa

Rabbit Anti-Human AMPK alpha-1 Antibody Bind with Human AMPK alpha-1 protein, His Tag



Measured by its binding ability in a indirect ELISA. Immobilized Human AMPK alpha-1 protein, His Tag (Cat. bs-41337P) at 2 µg/mL (100 µL/well) can bind Rabbit Anti-Human AMPK alpha-1 Antibody, the EC50 is 13.01 ng/mL.

Rabbit Anti-Human AMPK alpha-1 Antibody Bind with Human AMPK alpha-1 protein, His Tag



Measured by its binding ability in a indirect ELISA. Immobilized Human AMPK alpha-1 protein, His Tag (Cat. bs-41337P) at 2 µg/mL (100 µL/well) can bind Rabbit Anti-Human AMPK alpha-1 Antibody, the EC50 is 133.7 ng/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.