

human CD123 Mouse mAb

human CD123 Mouse mAb

Catalog # AP94211

Product Information

Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Physical State	Liquid
Isotype	Mouse IgG1, k
Purity	affinity purified by Protein G
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is an interleukin 3 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2 receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y pseudoautosomal region on chromosomes X or Y. Alternatively spliced transcript variants encoding distinct isoforms have been found. [provided by RefSeq, Jun 2012]

Additional Information


Dilution	Flow-Cyt=1:50-100
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background


This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images

scatter diagram showing peripheral blood lymphocytes stained with CD123. The cells were incubated with the



antibody (AP94211) for 30 min at 22°C. The secondary antibody used for 40 min at room temperature. Acquisition of >10,000 events was performed.



scatter diagram showing peripheral blood lymphocytes stained with CD123. The cells were incubated with the antibody (AP94211) for 30 min at 22°C. The secondary antibody used for 40 min at room temperature. Acquisition of >10,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.