

AEBP1 Rabbit pAb

AEBP1 Rabbit pAb
Catalog # AP94248

Product Information

Application	IHC-P, IHC-F, IF
Reactivity	Mouse, Rat
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	128 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse AEBP1
Epitope Specificity	951-1050/1128
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Isoform 1: Secreted. Isoform 2: Cytoplasm (Probable). Nucleus (Probable).
SIMILARITY	Belongs to the peptidase M14 family. Contains 1 F5/8 type C domain.
SUBUNIT	Interacts with GNG5, NFKBIA, MAPK1, MAPK3 and PTEN. May interact with calmodulin. Binds to DNA in vitro.
Post-translational modifications	Phosphorylated by MAPK1 in vitro.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of carboxypeptidase A protein family. The encoded protein may function as a transcriptional repressor and play a role in adipogenesis and smooth muscle cell differentiation. Studies in mice suggest that this gene functions in wound healing and abdominal wall development. Overexpression of this gene is associated with glioblastoma. [provided by RefSeq, May 2013].

Additional Information


Target/Specificity	Expressed in osteoblast and visceral fat.
Dilution	IHC-P=1:400-800,IHC-F=1:400-800,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This gene encodes a member of carboxypeptidase A protein family. The encoded protein may function as a

transcriptional repressor and play a role in adipogenesis and smooth muscle cell differentiation. Studies in mice suggest that this gene functions in wound healing and abdominal wall development. Overexpression of this gene is associated with glioblastoma. [provided by RefSeq, May 2013].

Images



Paraformaldehyde-fixed, paraffin embedded (rat small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (AEBP1) Polyclonal Antibody, Unconjugated (AP94248) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.