

THRB Rabbit pAb

THRB Rabbit pAb
Catalog # AP94251

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P10828
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52788
Physical State	Liquid
Immunogen	Recombinant human THRB protein
Epitope Specificity	209-461/461
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
DISEASE	Defects in THRB are the cause of generalized thyroid hormone resistance (GTHR) [MIM:188570, 274300]. GTHR is transmitted as an autosomal dominant trait, but an autosomal recessive form also exists. The disease is characterized by goiter, abnormal mental functions, increased susceptibility to infections, abnormal growth and bone maturation, tachycardia and deafness. Affected individuals may also have attention deficit-hyperactivity disorders (ADHD) and language difficulties. GTHR patients also have high levels of circulating thyroid hormones (T3-T4), with normal or slightly elevated thyroid stimulating hormone (TSH).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Thyroid hormone receptors (TRs) are ligand-dependent transcription factors that mediate the biological activities of thyroid hormone (T3). Thyroid hormone receptor b2 (TRb2) is a high affinity receptor for triiodothyronine which belongs to the nuclear hormone receptor family and the NR1 subfamily. It is composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal steroid-binding domain. Defects in the receptor result in generalized thyroid hormone resistance (GTHR). GTHR is transmitted as an autosomal dominant trait, but an autosomal recessive form also exists. The disease is characterized by goiter, abnormal mental functions, increased susceptibility to infections, abnormal growth and bone maturation, tachycardia and deafness. GTHR patients also have high levels of circulating thyroid hormones (T3-T4), with normal or slightly elevated thyroid stimulating hormone.

Additional Information

Gene ID 7068

Other Names	Thyroid hormone receptor beta, Nuclear receptor subfamily 1 group A member 2, c-erbA-2, c-erbA-beta, THRB, ERBA2, NR1A2, THR1
Dilution	IHC-P=1:200-800,IHC-F=1:400-800,IF=1:100-500,Flow-Cyt=1ug/Test,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	THRB
Synonyms	ERBA2, NR1A2, THR1
Function	Nuclear hormone receptor that can act as a repressor or activator of transcription. High affinity receptor for thyroid hormones, including triiodothyronine and thyroxine.
Cellular Location	Nucleus.

Background



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images




Blank control(black line):A431.
 Primary Antibody (green line): Rabbit Anti-THRB antibody (AP94251)
 Dilution:1ug/Test;
 Secondary Antibody(white blue line): Goat anti-rabbit IgG-AF488
 Dilution: 0.5ug/Test.
 Isotype control(orange line): Normal Rabbit IgG
 Protocol
 The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.


Measured by its binding ability in a indirect ELISA.
 Immobilized Human THRB protein, His Tag (Cat. AP94251) at 2 µg/mL (100 µL/well) can bind Rabbit Anti-Human THRB Antibody, the EC50 is 942.9 ng/mL.




Paraformaldehyde-fixed, paraffin embedded (human thyroid gland); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (THRB) Polyclonal Antibody, Unconjugated (AP94251) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Measured by its binding ability in a indirect ELISA. Immobilized Human THRB protein, His Tag (Cat. bs-41495P) at 2 µg/mL (100 µL/well) can bind Rabbit Anti-Human THRB Antibody, the EC50 is 133.9 ng/mL.




Measured by its binding ability in a indirect ELISA. Immobilized Human THRB protein, His Tag (Cat. bs-41495P) at 2 µg/mL (100 µL/well) can bind Rabbit Anti-Human THRB Antibody, the EC50 is 942.9 ng/mL.



Blank control(black line):A431. Primary Antibody (green line): Rabbit Anti-THRB antibody (AP94251) Dilution:1ug/Test; Secondary Antibody(white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control(orange line): Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control(black line):A431. Primary Antibody (green line): Rabbit Anti-THRB antibody (AP94251)
Dilution:1ug/Test; Secondary Antibody(white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control(orange line): Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control(black line):Hela. Primary Antibody (green line): Rabbit Anti-THRB antibody (AP94251)
Dilution:1ug/Test; Secondary Antibody(white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control(orange line): Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.