

FRA10AC1 Rabbit pAb

FRA10AC1 Rabbit pAb

Catalog # AP94389

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	Human
Predicted	Rat, Pig, Dog, Mouse, Rabbit, Chicken, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C10orf4
Epitope Specificity	101-200/315
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nuclear.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a nuclear phosphoprotein of unknown function. This gene contains a tandem CGG repeat region within a CpG island that normally consists of 8-14 repeats but can expand to over 200 repeats. The repeat region is within the 5' UTR of some transcript variants, but is intronic to another variant. The expanded repeat allele is a fragile site and becomes hypermethylated, causing a reduction in gene expression. A disease phenotype has not been associated with expanded alleles. This gene is found within the rare FRA10A folate-sensitive fragile site. [provided by RefSeq, Dec 2016]


Additional Information

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:


293T(Human) Cell Lysate at 40 ug

Primary: Anti-C10orf4 (AP94389) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 38 kD

Observed band size: 38 kD



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C10orf4) Polyclonal Antibody, Unconjugated (AP94389) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.