

SMARCA2/BRM Recombinant Rabbit mAb

SMARCA2/BRM Recombinant Rabbit mAb

Catalog # AP94391

Product Information

Application	IHC-P, IHC-F, IF
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Recombinant
Physical State	Liquid
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	A transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. SMARCA2 / BRM belongs to the SNF2/RAD54 helicase family, is a homologue of the <i>Saccharomyces cerevisiae</i> SWI2/SNF2 and <i>Drosophila</i> brahma proteins. It contains a methyl lysine containing bromo domain and an HSA domain. The yeast protein SNF2, also known as SWI2, is involved in transcriptional activation of numerous genes. It contains a domain that is highly conserved among several known helicases and is required for transcriptional activity. SNF2/SWI2 is highly homologous to the <i>Drosophila</i> protein 'brahma' (brm). Although the 2 proteins show nuclear localization during interphase, they are excluded from the condensed chromosomes during mitosis. They found that the level of BRM, but not BRG1, was strongly reduced during mitosis. Phosphorylation of hbrm and BRG1 did not disrupt their association with SNF5 but correlated with a decreased affinity for the nuclear structure in early M phase.

Additional Information

Dilution	IHC-P=1:50-200,IHC-F=1:50-200,IF=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.