

TROP2 Recombinant Rabbit mAb

TROP2 Recombinant Rabbit mAb

Catalog # AP94410

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	Human
Host	Rabbit
Clonality	Recombinant
Physical State	Liquid
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein.
DISEASE	Corneal dystrophy, gelatinous drop-like (GDLD) [MIM:204870]: A form of lattice corneal dystrophy, a class of inherited stromal amyloidoses characterized by pathognomonic branching lattice figures in the cornea. GDLD is an autosomal recessive disorder characterized by severe corneal amyloidosis leading to blindness. Clinical manifestations, which appear in the first decade of life, include blurred vision, photophobia, and foreign-body sensation. By the third decade, raised, yellowish-gray, gelatinous masses severely impair visual acuity. Note=The disease is caused by mutations affecting the gene represented in this entry.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This intronless gene encodes a carcinoma-associated antigen. This antigen is a cell surface receptor that transduces calcium signals. Mutations of this gene have been associated with gelatinous drop-like corneal dystrophy.[provided by RefSeq, Dec 2009]


Additional Information

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



25 ug total protein per lane of various lysates (see on figure) probed with TROP2 monoclonal antibody, unconjugated (AP94410) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Breast; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with TROP2 Monoclonal Antibody, Unconjugated (AP94410) at 1:200 overnight at 4°C, followed by conjugation to the AP94410-HRP and DAB (C-0010) staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.