

SLC22A12 Rabbit pAb

SLC22A12 Rabbit pAb

Catalog # AP94412

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q96S37
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59630
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SLC22A12
Epitope Specificity	101-200/553
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane, Multi-pass membrane protein.
DISEASE	Hypouricemia renal 1 (RHUC1) [MIM:220150]: A disorder characterized by impaired uric acid reabsorption at the apical membrane of proximal renal tubule cells, and high urinary urate excretion. Patients often appear asymptomatic, but may be subject to exercise-induced acute renal failure, chronic renal dysfunction and nephrolithiasis. Note=The disease is caused by mutations affecting the gene represented in this entry.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is involved in the sodium-independent transport and excretion of organic anions, some of which are potentially toxic. The encoded protein is an integral membrane protein and is found mainly in the kidney and in the placenta, where it may act to prevent potentially harmful organic anions from reaching the fetus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015]

Additional Information

Gene ID	116085
Other Names	Solute carrier family 22 member 12, Organic anion transporter 4-like protein, Renal-specific transporter, RST {ECO:0000303 Ref.2}, Urate anion exchanger 1, URAT1, Urate:anion antiporter SLC22A12, SLC22A12 (HGNC:17989)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	SLC22A12 (HGNC:17989)
Function	Electroneutral antiporter that translocates urate across the apical membrane of proximal tubular cells in exchange for monovalent organic or inorganic anions (PubMed: 12024214 , PubMed: 22194875 , PubMed: 35144162 , PubMed: 35462902). Involved in renal reabsorption of urate and helps maintaining blood levels of uric acid (PubMed: 12024214 , PubMed: 22194875). Mediates urate uptake by an exchange with organic anions such as (S)-lactate and nicotinate, and inorganic anion Cl(-) (PubMed: 12024214). Other inorganic anions such as Br(-), I(-) and NO3(-) may also act as counteranions that exchange for urate (PubMed: 12024214). Also mediates orotate tubular uptake coupled with nicotinate efflux and to a lesser extent with lactate efflux, therefore displaying a potential role in orotate renal reabsorption (PubMed: 21350910). Orotate transport is Cl(-)-dependent (PubMed: 21350910).
Cellular Location	Apical cell membrane; Multi-pass membrane protein
Tissue Location	Detected in kidney (at protein level). Detected in fetal and adult kidney. Detected in epithelial cells of proximal tubules in renal cortex.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: HeLa Cell (Human) Lysate at 40 ug
Primary: Anti-SLC22A12 (AP94412) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 61 kD
Observed band size: 61 kD



Tissue/cell: human kidney tissue; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-SLC22A12 Polyclonal Antibody, Unconjugated(AP94412) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.