

phospho-ERK1/2 (Thr202 + Tyr204) Rabbit pAb

phospho-ERK1/2 (Thr202 + Tyr204) Rabbit pAb

Catalog # AP94421

Product Information


Reactivity	Mouse
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated Synthesised phosphopeptide derived from mouse p44/42 MAPK around the phosphorylation site of Thr202/Tyr204
Epitope Specificity	FL(p-T)E(p-Y)VA
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm, cytoskeleton, spindle. Nucleus. Cytoplasm, cytoskeleton, centrosome. Cytoplasm. Note=Associated with the spindle duringprometaphase and metaphase. PEA15-binding andphosphorylated DAPK1 promote its cytoplasmic retention.Phosphorylation at Ser-244 and Ser-246 as well asautophosphorylation at Thr-188 promote nuclear localization.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a member of the MAPkinase family. MAP kinases, also known as extracellularsignal-regulated kinases (ERKs), act in a signaling cascade thatregulates various cellular processes such as proliferation,differentiation, and cell cycle progression in response to avariety of extracellular signals. This kinase is activated byupstream kinases, resulting in its translocation to the nucleuswhere it phosphorylates nuclear targets. Alternatively splicedtranscript variants encoding different protein isoforms have beendescribed. [provided by RefSeq, Jul 2008].

Additional Information


Dilution	ICC/IF=1:100-500,Flow-Cyt=1:50-100
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background


This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.




Blank control (blue line): U251 (fixed with 2% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min at room temperature).
 Primary Antibody (green line): Rabbit Anti-phospho-ERK12 (Thr202 + Tyr204) antibody (AP94421), Dilution: 3 μ g / 10^6 cells;
 Isotype Control Antibody (orange line): Rabbit IgG .
 Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE, Dilution: 1 μ g / test.



Tissue/cell: HUVEC cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (phospho-ERK1/2 (Thr202 + Tyr204)) polyclonal Antibody, Unconjugated (AP94421) 1:100, 90 minutes at 37°C; followed by a FITC conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.





Paraformaldehyde-fixed, paraffin embedded (rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30 min; Antibody incubation with (phospho-ERK12 (Thr202 + Tyr204)) Polyclonal Antibody, Unconjugated (AP94421) at 1:2000 overnight at 4°C, followed by operating according to SP Kit (Rabbit) (sp-0023) instructions and DAB staining.



Blank control (blue line): U251 (fixed with 2% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min at room temperature).
 Primary Antibody (green line): Rabbit Anti-phospho-ERK12 (Thr202 + Tyr204) antibody (AP94421), Dilution: 3 μ g / 10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE, Dilution: 1 μ g / test.

Tissue/cell: Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (phospho-ERK1/2 (Thr202 + Tyr204)) polyclonal Antibody, Unconjugated (AP94421) 1:100, 90 minutes at 37°C; followed by a FITC conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Tissue/cell: HUVEC cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (phospho-ERK1/2 (Thr202 + Tyr204)) polyclonal Antibody, Unconjugated (AP94421) 1:100, 90 minutes at 37°C; followed by a FITC conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.