

NUP62 Recombinant Mouse mAb

NUP62 Recombinant Mouse mAb

Catalog # AP94427

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Recombinant
Physical State	Liquid
Isotype	IgG1, Kappa
Purity	affinity purified by Protein G
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus, nuclear pore complex. Cytoplasm, cytoskeleton, spindle pole. Note=Central region of the nuclear pore, within the transporter. During mitotic cell division, it associates with the poles of the mitotic spindle.
SIMILARITY	Belongs to the nucleoporin NSP1/NUP62 family.
SUBUNIT	Component of the p62 complex, a complex at least composed of NUP62, NUP54, and NUPL1 (By similarity). Interacts with C11orf73/Hikeshi.
Post-translational modifications	O-glycosylated. Contains about 10 N-acetylglucosamine side chain sites predicted for the entire protein, amongst which only one in the C-terminal.
DISEASE	Infantile striatonigral degeneration (SNDI) [MIM:271930]: Neurological disorder characterized by symmetrical degeneration of the caudate nucleus, putamen, and occasionally the globus pallidus, with little involvement of the rest of the brain. The clinical features include developmental regression, choreoathetosis, dystonia, spasticity, dysphagia, failure to thrive, nystagmus, optic atrophy, and mental retardation. Note=The disease is caused by mutations affecting the gene represented in this entry.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The nuclear pore complex is a massive structure that extends across the nuclear envelope, forming a gateway that regulates the flow of macromolecules between the nucleus and the cytoplasm. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. The protein encoded by this gene is a member of the FG-repeat containing nucleoporins and is localized to the nuclear pore central plug. This protein associates with the importin alpha/beta complex which is involved in the import of proteins containing nuclear localization signals. Multiple transcript variants of this gene encode a single protein isoform. [provided by RefSeq].

Additional Information

Dilution	WB=1:500-2000,IHC-P=1:50-200,IHC-F=1:50-200,IF=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Background

The nuclear pore complex is a massive structure that extends across the nuclear envelope, forming a gateway that regulates the flow of macromolecules between the nucleus and the cytoplasm. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. The protein encoded by this gene is a member of the FG-repeat containing nucleoporins and is localized to the nuclear pore central plug. This protein associates with the importin alpha/beta complex which is involved in the import of proteins containing nuclear localization signals. Multiple transcript variants of this gene encode a single protein isoform. [provided by RefSeq].

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.