

SEPT2 Recombinant Mouse mAb

SEPT2 Recombinant Mouse mAb

Catalog # AP94498

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	Human
Predicted	Mouse, Rat
Host	Mouse
Clonality	Recombinant
Physical State	Liquid
Isotype	IgG1, Kappa
Purity	affinity purified by Protein G
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, spindle. Chromosome, centromere, kinetochore. Cleavage furrow. Midbody. Cytoplasm, cell cortex.
SIMILARITY	Belongs to the septin family.
SUBUNIT	Septins polymerize into heterooligomeric protein complexes that form filaments, and associate with cellular membranes, actin filaments and microtubules.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The septins are a family of cytoskeletal GTPases that play an essential role in cytokinesis in mammalian cells. Septin 2 (Nedd5) is a mammalian septin known to associate with actin based structures such as the contractile ring and stress fibers. It is involved in cytokinesis and may assemble into a multicomponent structure.

Additional Information

Target/Specificity	Widely expressed. Up-regulated in liver cancer.
Dilution	WB=1:500-1000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

The septins are a family of cytoskeletal GTPases that play an essential role in cytokinesis in mammalian cells. Septin 2 (Nedd5) is a mammalian septin known to associate with actin based structures such as the contractile ring and stress fibers. It is involved in cytokinesis and may assemble into a multicomponent structure.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.