

ATP5A1 Recombinant Mouse mAb

ATP5A1 Recombinant Mouse mAb

Catalog # AP94569

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Recombinant
Physical State	Liquid
Isotype	IgG2b/Kappa
Purity	affinity purified by Protein G
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Mitochondrion. Mitochondrion inner membrane.
SIMILARITY	Belongs to the eukaryotic ATPase subunit F6 family.
SUBUNIT	F-type ATPases have 2 components, CF(1) - the catalytic core - and CF(0) - the membrane proton channel. CF(0) seems to have nine subunits: a, b, c, d, e, f, g, F6 and 8 (or A6L).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	ATP5J (ATP synthase, H ⁺ transporting, mitochondrial F0 complex, subunit F6) is a multisubunit membrane-bound enzyme complex consisting of an F0 segment embedded in the membrane and an F1 segment attached to the F0. It is also a component of mitochondrial ATP synthase which is required for the interactions of the catalytic and proton-translocating segments. Human ATP5J shares 72% sequence identity with rat ATP5J. This signal peptide is rich in basic amino acids, devoid of acidic amino acids, and amphiphilic, which allows it to be water-soluble yet capable of passage through the phospholipid membrane bilayers. Moreover, it is circulating and functions as an endogenous vasoconstrictor by inhibiting cytosolic phospholipase A2.

Additional Information

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:50-200,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

ATP5J (ATP synthase, H⁺ transporting, mitochondrial F0 complex, subunit F6) is a multisubunit membrane-bound enzyme complex consisting of an F0 segment embedded in the membrane and an F1 segment attached to the F0. It is also a component of mitochondrial ATP synthase which is required for the

interactions of the catalytic and proton-translocating segments. Human ATP5J shares 72% sequence identity with rat ATP5J. This signal peptide is rich in basic amino acids, devoid of acidic amino acids, and amphiphilic, which allows it to be water-soluble yet capable of passage through the phospholipid membrane bilayers. Moreover, it is circulating and functions as an endogenous vasoconstrictor by inhibiting cytosolic phospholipase A2.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.