

VIPR2 Rabbit pAb

VIPR2 Rabbit pAb
Catalog # AP94599

Product Information

Application	WB, IHC-P, IHC-F, IF
Reactivity	Human
Predicted	Rat, Mouse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from rat VIP receptor II
Epitope Specificity	81-170/437
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Multi-pass membrane protein.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Pituitary adenylate cyclase-activating polypeptide (PACAP) is a neuropeptide belonging to the vasoactive intestinal polypeptide/glucagon/ secretin family. It is widely distributed in the body, and a variety of biological actions have been reported. Recent studies have shown that there is a family of PACAP receptors (PACAPRs), and two members of this family have been identified. Mouse PACAPR-3 is a protein of 437 amino acids that has 50% and 51% identity with rat PACAP type I and type II receptors, respectively. It binds to vasoactive intestinal polypeptide as well as PACAP-38 and -27, with a slightly higher affinity for PACAP-38, PACAPR-3 mRNA is expressed at high levels in MIN6, at moderate levels in pancreatic islets and other insulin-secreting cell lines, HIT-T15 and RINm5F, as well as in the lung, brain, stomach, and colon, and at low levels in the heart. PACAPR-3 participates in the regulation of insulin secretion. Insulin secretion from MIN6 cells is significantly stimulated by PACAP-38.


Additional Information

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=2ug/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Blank control:Hela.

Primary Antibody (green line): Rabbit Anti-VPAC2R antibody (AP94599)

Dilution: 2 µg /10⁶ cells;


Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 2 µg /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.




Sample: A431 Cell Lysate at 40 µg

Primary: Anti- VPAC2R (AP94599) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 48 kD

Observed band size: 63 kD



Tissue/cell: human lung carcinoma; 4%

Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3%

Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-VIP receptor 2/VPAC2 Polyclonal

Antibody, Unconjugated(AP94599) 1:200, overnight at 4°C, followed by conjugation to the secondary

antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.