

LDL Rabbit pAb

LDL Rabbit pAb
Catalog # AP94684

Product Information

Application	IHC-P, IHC-F, IF
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	Lipoprotein, low density from human plasma
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	LDL (low-density lipoprotein) is a type of lipoprotein that transports cholesterol and triglycerides from the liver to peripheral tissues. LDL enables fats and cholesterol to move within the water based solution of the blood stream. LDL also regulates cholesterol synthesis at these sites. The low density lipoprotein (LDL) receptor system coordinates the metabolism of cholesterol, an essential component of the plasma membrane of all mammalian cells. Study of this system has led to an enhanced understanding of the cellular basis of cholesterol homeostasis. It has also brought into focus an important mechanism of metabolic regulation--the process of receptor-mediated endocytosis (1). Data suggest that the juxtamembranous region of the cytoplasmic domain participates in protein:protein interactions that allow the low density lipoprotein receptor to cluster in coated pits (2). It has been shown that the family of LDL receptors may serve as viral receptors. Endocytosis of the Flaviviridae viruses, hepatitis C virus, GB virus C/hepatitis G virus, and bovine viral diarrheal virus (BVDV) was shown to be mediated by LDL receptors on cultured cells.


Additional Information

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=3ug/test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Blank control: A549.

Primary Antibody (green line): Rabbit Anti-LDL antibody (AP94684)

Dilution: 3 µg /10⁶ cells;


Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 µg /test.

Protocol

The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Tissue/cell: Human lung cancer; 4%

Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling

bathing for 15min; Block endogenous peroxidase by 3%

Hydrogen peroxide for 30min; Blocking buffer (normal


goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-LDL Polyclonal Antibody,

Unconjugated(AP94684) 1:200, overnight at 4°C, followed

by conjugation to the secondary antibody(SP-0023) and

DAB(C-0010) staining



Blank control: A549. Primary Antibody (green line): Rabbit

Anti-LDL antibody (AP94684) Dilution: 3 µg /10⁶ cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE Dilution: 1

µg /test. Protocol The cells were incubated in 5%BSA to

block non-specific protein-protein interactions for 30 min

at room temperature .Cells stained with Primary

Antibody for 30 min at room temperature. The secondary

antibody used for 40 min at room temperature.

Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.