

# EAR1 Rabbit pAb

EAR1 Rabbit pAb  
Catalog # AP94717

## Product Information

---

|                                |                                                                                                                                                                                                                                            |
|--------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Application</b>             | WB, IHC-P, IHC-F, IF                                                                                                                                                                                                                       |
| <b>Reactivity</b>              | Rat, Human, Mouse                                                                                                                                                                                                                          |
| <b>Host</b>                    | Rabbit                                                                                                                                                                                                                                     |
| <b>Clonality</b>               | Polyclonal                                                                                                                                                                                                                                 |
| <b>Physical State</b>          | Liquid                                                                                                                                                                                                                                     |
| <b>Immunogen</b>               | KLH conjugated synthetic peptide derived from mouse Eosinophil cationic protein 1                                                                                                                                                          |
| <b>Epitope Specificity</b>     | 65-155/155                                                                                                                                                                                                                                 |
| <b>Isotype</b>                 | IgG                                                                                                                                                                                                                                        |
| <b>Purity</b>                  | affinity purified by Protein A                                                                                                                                                                                                             |
| <b>Buffer</b>                  | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.                                                                                                                                                                          |
| <b>SUBCELLULAR LOCATION</b>    | Secreted. Note=Located in the matrix of eosinophil large specific granule, which are released following activation by an immune stimulus.                                                                                                  |
| <b>Important Note</b>          | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.                                                                                                                  |
| <b>Background Descriptions</b> | Eosinophil derived neurotoxin (EDN) is a protein belonging to the ribonuclease (RNase) A superfamily. It has recently been found to have antiviral activity against respiratory syncytial virus and human immunodeficiency virus in vitro. |

## Additional Information

---

|                 |                                                                                                                                                                                                 |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Dilution</b> | WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1 ug /Test                                                                                                                  |
| <b>Storage</b>  | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

## Background


---

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

## Images

---

Blank control (Black line): Molt4 (Black).  
Primary Antibody (green line): Rabbit Anti-EAR1 antibody (AP94717)




Dilution: 1  $\mu\text{g}$  /  $10^6$  cells;  
 Isotype Control Antibody (orange line): Rabbit IgG .  
 Secondary Antibody (white blue line): Goat anti-rabbit  
 IgG-AF647

Dilution: 1  $\mu\text{g}$  /test.

#### Protocol

The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



#### Sample:

Lane 1: Mouse Spleen tissue lysates


Lane 2: Mouse Bone tissue lysates

Primary: Anti-EAR1 (AP94717) at 1/200 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 14 kDa

Observed band size: 17 kDa



Paraformaldehyde-fixed, paraffin embedded (rat pancreas tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EAR1) Polyclonal Antibody, Unconjugated (AP94717) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.