

SPON1 Rabbit pAb

SPON1 Rabbit pAb
Catalog # AP94764

Product Information

Application	IHC-P, IHC-F, IF
Reactivity	Rat, Mouse
Predicted	Pig, Dog, Human, Rabbit
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SPON1/F spondin
Epitope Specificity	741-807/807
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted, extracellular space, extracellular matrix.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	SPON1 is a member of a subgroup of the thrombospondin type 1 (TSR) class molecules, defined by two domains of homology, the FS1/FS2 and TSR domains. The TSRs of SPON1 proteins are typical of class 2 TSRs. SPON1, which is similar to thrombospondin, is a extracellular matrix attached molecule that promotes neurite outgrowth and inhibits angiogenesis. Analysis of gain and loss of function experiments reveal that SPON1 is required for accurate pathfinding of embryonic axons, and plays a dual role in patterning axonal trajectories. It promotes the outgrowth of commissural and inhibits the outgrowth of motor axons, and has also been implicated in inflammatory processes in the nervous system.


Additional Information

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SPON1) Polyclonal Antibody, Unconjugated (AP94764) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.