10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MUC6 Mouse mAb

MUC6 Mouse mAb
Catalog # AP94792

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q6W4X9
Reactivity	Rat, Human, Mouse
Host	Mouse
Clonality	Monoclonal
Calculated MW	257051
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MUC6
Isotype	IgG1, Kappa
Purity	affinity purified by Protein A
Buffer	Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02% Proclin300.
SUBCELLULAR LOCATION	Secreted.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins produced by many epithelial tissues. The protein encoded by this gene is secreted and forms an insoluble mucous barrier that protects the gut lumen. [provided by RefSeq, Dec 2016]

Additional Information

Gene ID	4588
Other Names	Mucin-6, MUC-6, Gastric mucin-6, MUC6
Dilution	IHC-P=1:100-500,IF=1:100-500,IHC-F=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	MUC6
Function	May provide a mechanism for modulation of the composition of the protective mucus layer related to acid secretion or the presence of bacteria and noxious agents in the lumen. Plays an important role in the cytoprotection of epithelial surfaces and are used as tumor markers in a

	variety of cancers. May play a role in epithelial organogenesis.
Cellular Location	Secreted.
Tissue Location	Expressed in the regenerative zone of gastric antrum, gastric body mucosa and gastric incisura mucosa. Expressed in the deeper mucous glands of gastric antrum. Overexpressed in Helicobacter pylori infected gastric epithelium. Highly expressed in duodenal Brunner's glands, gall bladder, seminal vesicle, pancreatic centroacinar cells and ducts, and periductal glands of the common bile duct.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (human stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (MUC6) Monoclonal Antibody, Unconjugated (AP94792) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructionsand DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.