10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Recombinant Anti-Cytokeratin 16 Rabbit mAb

Catalog # AP94939

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	P08779
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	51268


Additional Information

Gene ID	3868
Other Names	KRT16A; Keratin, type I cytoskeletal 16; Cytokeratin-16; CK-16; Keratin-16; K16
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human Cytokeratin 16 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	KRT16 (HGNC:6423)
Synonyms	KRT16A
Function	Structural component of intermediate filaments in epithelial keratinocytes. Forms heteropolymers with type II keratins KRT6A and KRT6B, assembling into keratin intermediate filament networks that contribute to cytoskeletal resilience of palm, sole, nail bed and oral mucosa (PubMed:7539673, PubMed:8595410). Regulates epidermal innate immunity and inflammatory signaling in response to barrier disruption. Required for normal palmoplantar epidermal differentiation and nail unit structure (By similarity).
Tissue Location	Expressed in the corneal epithelium (at protein level).


Images



Western blot analysis of Cytokeratin 16 expression in HaCat (A) whole cell lysates. (Predicted band size: 51 kD; Observed band size: 51 kD)



Immunohistochemical analysis of Cytokeratin 16 staining in human skin formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Cytokeratin 16 staining in HaCat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.