

Recombinant Anti-Caspase 10 Rabbit mAb

Catalog # AP94972

Product Information

Application	WB, IHC, FC, IP, IF/IC
Primary Accession	Q92851
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal
Calculated MW	58951


Additional Information

Gene ID	843
Other Names	MCH4; Caspase-10; CASP-10; Apoptotic protease Mch-4; FAS-associated death domain protein interleukin-1B-converting enzyme 2; FLICE2; ICE-like apoptotic protease 4
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human Caspase 10 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	CASP10
Synonyms	MCH4
Function	Involved in the activation cascade of caspases responsible for apoptosis execution. Recruited to both Fas- and TNFR-1 receptors in a FADD dependent manner. May participate in the granzyme B apoptotic pathways. Cleaves and activates effector caspases CASP3, CASP4, CASP6, CASP7, CASP8 and CASP9. Hydrolyzes the small- molecule substrates, Tyr- Val-Ala-Asp- -AMC and Asp-Glu-Val-Asp- -AMC.
Tissue Location	Detectable in most tissues. Lowest expression is seen in brain, kidney, prostate, testis and colon

Images



Western blot analysis of Caspase 10 expression in HeLa (A) whole cell lysates. (Predicted band size: 59 kD; Observed band size: 70 kD)



Immunofluorescent analysis of Caspase 10 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.