

Recombinant Anti-CD120b Rabbit mAb

Catalog # AP95032

Product Information

Application	WB, IHC, FC, IP, IF/IC
Primary Accession	P20333
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	48291


Additional Information

Gene ID	7133
Other Names	TNFR2; TNFRSF1B; Tumor necrosis factor receptor superfamily member 1B; Tumor necrosis factor receptor 2; TNF-R2; Tumor necrosis factor receptor type II; TNF-RII; TNFR-II; p75; p80 TNF-alpha receptor; CD120b; Etanercept
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human CD120b protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C. Stable for 12 months from date of receipt


Protein Information

Name	TNFRSF1B
Synonyms	TNFR2, TNFR
Function	Receptor with high affinity for TNFSF2/TNF and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF. Isoform 2 blocks TNF-induced apoptosis, which suggests that it regulates TNF function by antagonizing its biological activity.
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Tumor necrosis factor-binding protein 2]: Secreted

Images



Western blot analysis of CD120b expression in Jurkat (A) whole cell lysates. (Predicted band size: 48 kD; Observed band size: 65 kD)



Immunofluorescent analysis of CD120b staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.