

Recombinant Anti-DR5 Rabbit mAb

Catalog # AP95054

Product Information

Application	WB, FC, IP
Primary Accession	O14763
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	47878

Additional Information


Gene ID	8795
Other Names	DR5; KILLER; TRAILR2; TRICK2; ZTNFR9; Tumor necrosis factor receptor superfamily member 10B; Death receptor 5; TNF-related apoptosis-inducing ligand receptor 2; TRAIL receptor 2; TRAIL-R2; CD262
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human DR5 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IP~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	TNFRSF10B
Synonyms	DR5, KILLER, TRAILR2, TRICK2, ZTNFR9
Function	Receptor for the cytotoxic ligand TNFSF10/TRAIL (PubMed: 10549288). The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and

throughout the intestinal tract; not detectable in brain

Images



Western blot analysis of DR5 expression in HeLa (A) whole cell lysates. (Predicted band size: 48 kD; Observed band size: 40, 48 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.