

Recombinant Anti-BUB1B Rabbit mAb

Catalog # AP95055

Product Information

Application	WB, IHC, FC, IP
Primary Accession	O60566
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	119545

Additional Information

Gene ID	701
Other Names	BUBR1; MAD3L; SSK1; Mitotic checkpoint serine/threonine-protein kinase BUB1 beta; MAD3/BUB1-related protein kinase; hBUBR1; Mitotic checkpoint kinase MAD3L; Protein SSK1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human BUB1B protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	BUB1B
Synonyms	BUBR1, MAD3L, SSK1
Function	Component of the mitotic checkpoint complex (MCC), which inhibits APC/C and delays chromosome segregation until all chromosomes are properly attached to the mitotic spindle (PubMed:27509861). The MCC, which consists of CDC20, MAD2L1/MAD2, BUB1B/BUBR1 and BUB3, is a key player of the spindle assembly checkpoint (PubMed:11702782, PubMed:14706340, PubMed:15020684, PubMed:27509861). The MCC-bound APC/C complex is inactive and this delays the metaphase/anaphase transition (PubMed:10477750, PubMed:27509861). When the spindle assembly checkpoint is silenced, the MCC-APC/C complex recruits the E2 enzyme UBCH10 and triggers the auto-ubiquitination of CDC20 in the MCC, thereby reactivating APC/C for transition into anaphase (PubMed:27509861). MCC also monitors kinetochore activities that depend on the kinetochore motor CENPE (PubMed:15020684). Required for kinetochore localization of CENPE

(PubMed:[15020684](#)). Negatively regulates PLK1 activity in interphase cells and suppresses centrosome amplification (PubMed:[19503101](#)). May play a role for tumor suppression (By similarity).


Cellular Location

Cytoplasm. Nucleus. Chromosome, centromere, kinetochore {ECO:0000250|UniProtKB:Q9Z1S0}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Cytoplasmic in interphase cells. Associates with the kinetochores in early prophase Kinetochore localization requires BUB1, PLK1 and KNL1

Tissue Location

Highly expressed in thymus followed by spleen. Preferentially expressed in tissues with a high mitotic index

Images



Western blot analysis of BUB1B expression in HepG2 (A) whole cell lysates. (Predicted band size: 120 kD; Observed band size: 120 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.