

Recombinant Anti-N-WASP Rabbit mAb

Catalog # AP95063

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	O00401
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal
Calculated MW	54827

Additional Information

Gene ID	8976
Other Names	Neural Wiskott-Aldrich syndrome protein; N-WASP
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human N-WASP protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	WASL
Function	Regulates actin polymerization by stimulating the actin- nucleating activity of the Arp2/3 complex (PubMed: 16767080 , PubMed: 19366662 , PubMed: 19487689 , PubMed: 22847007 , PubMed: 22921828 , PubMed: 9422512). Involved in various processes, such as mitosis and cytokinesis, via its role in the regulation of actin polymerization (PubMed: 19366662 , PubMed: 19487689 , PubMed: 22847007 , PubMed: 22921828 , PubMed: 9422512). Together with CDC42, involved in the extension and maintenance of the formation of thin, actin-rich surface projections called filopodia (PubMed: 9422512). In addition to its role in the cytoplasm, also plays a role in the nucleus by regulating gene transcription, probably by promoting nuclear actin polymerization (PubMed: 16767080). Binds to HSF1/HSTF1 and forms a complex on heat shock promoter elements (HSE) that negatively regulates HSP90 expression (By similarity). Plays a role in dendrite spine morphogenesis (By similarity). Decreasing levels of DNMBP (using antisense RNA) alters apical junction morphology in cultured enterocytes, junctions curve instead of being nearly linear (PubMed: 19767742).


Cellular Location

Cytoplasm, cytoskeleton. Nucleus Cytoplasm {ECO:0000250|UniProtKB:Q91YD9}. Note=Preferentially localized in the cytoplasm when phosphorylated and in the nucleus when unphosphorylated (By similarity). Exported from the nucleus by an nuclear export signal (NES)-dependent mechanism to the cytoplasm (By similarity). {ECO:0000250|UniProtKB:Q91YD9}

Images



Western blot analysis of N-WASP expression in HeLa (A) whole cell lysates. (Predicted band size: 55 kD; Observed band size: 60 kD)



Immunofluorescent analysis of N-WASP staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.