

Recombinant Anti-GHR Rabbit mAb

Catalog # AP95076

Product Information


Application	WB, FC, IF/IC
Primary Accession	P10912
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	71500

Additional Information


Gene ID	2690
Other Names	Growth hormone receptor; GH receptor; Somatotropin receptor
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human GHR protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	GHR
Function	Receptor for pituitary gland growth hormone (GH1) involved in regulating postnatal body growth (PubMed: 1549776 , PubMed: 2825030 , PubMed: 8943276). On ligand binding, couples to the JAK2/STAT5 pathway (PubMed: 1549776 , PubMed: 15690087 , PubMed: 2825030 , PubMed: 8943276).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Note=On growth hormone binding, GHR is ubiquitinated, internalized, down-regulated and transported into a degradative or non-degradative pathway {ECO:0000250 UniProtKB:P19941} [Growth hormone-binding protein]: Secreted. Note=Complexed to a substantial fraction of circulating GH.
Tissue Location	Expressed in various tissues with high expression in liver and skeletal muscle. [Isoform 4]: Predominantly expressed in kidney, bladder, adrenal gland and brain stem (PubMed:1569971). Highly expressed in placental villi (PubMed:1569971, PubMed:8360189)



Western blot analysis of GHR expression in MCF7 (A) whole cell lysates. (Predicted band size: 72 kD; Observed band size: 140 kD)



Immunofluorescent analysis of GHR staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.