10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Recombinant Anti-ULK3 Rabbit mAb

Catalog # AP95086

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	Q6PHR2
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	53444

Additional Information


Gene ID	25989
Other Names	Serine/threonine-protein kinase ULK3; Unc-51-like kinase 3
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human ULK3 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	ULK3 {ECO:0000303|PubMed:19878745, ECO:0000312|HGNC:HGNC:19703}
Function	Serine/threonine protein kinase that acts as a regulator of smoothened signaling and autophagy (PubMed:19279323, PubMed:19878745, PubMed:20643644, PubMed:31279575). Acts as a positive regulator of smoothened signaling in the presence of hedgehog (DHH, IHH or SHH) ligand: autophosphorylates and mediates phosphorylation of full-length GLI2, promoting subsequent phosphorylation by CDK1 and GLI2 dissociation from SUFU and translocation to the nucleus (PubMed:19878745, PubMed:20643644, PubMed:31279575). Also phosphorylates as GLI1 and GLI3, although less efficiently (PubMed:19878745, PubMed:20643644). In the absence of hedgehog ligand, interacts with SUFU, thereby inactivating the protein kinase activity and preventing phosphorylation of GLI proteins (GLI1, GLI2 and/or GLI3) (PubMed:20643644). Also acts as a regulator of autophagy: following cellular senescence, able to induce autophagy (PubMed:19279323).
Cellular Location	Cytoplasm. Note=Localizes to pre-autophagosomal structure during cellular senescence.

Tissue Location	Widely expressed. Highest levels observed in fetal brain. In adult tissues, high levels in brain, liver and kidney, moderate levels in testis and adrenal gland and low levels in heart, lung, stomach, thymus, prostate and placenta. In the brain, highest expression in the hippocampus, high levels also detected in the cerebellum, olfactory bulb and optic nerve. In the central nervous system, lowest levels in the spinal cord

Images



Western blot analysis of ULK3 expression in HEK293T (A) whole cell lysates. (Predicted band size: 53 kD; Observed band size: 53 kD)



Immunofluorescent analysis of ULK3 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.