

Recombinant Anti-CD82 Rabbit mAb

Catalog # AP95090

Product Information

Application	WB, IHC, FC
Primary Accession	P27701
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	29626

Additional Information

Gene ID	3732
Other Names	KAI1; SAR2; ST6; TSPAN27; CD82 antigen; C33 antigen; IA4; Inducible membrane protein R2; Metastasis suppressor Kangai-1; Suppressor of tumorigenicity 6 protein; Tetraspanin-27; Tspan-27; CD82
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human CD82 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CD82
Synonyms	KAI1, SAR2, ST6, TSPAN27
Function	Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling (PubMed: 19497983). Participates thereby in diverse biological functions such as cell signal transduction, adhesion, migration and protein trafficking. Acts as a attenuator of EGF signaling, facilitating ligand-induced endocytosis of the receptor and its subsequent desensitization (PubMed: 10985391 , PubMed: 35538033). Mechanistically, modulates ligand- induced ubiquitination and trafficking of EGFR via E3 ligase CBL phosphorylation by PKC (PubMed: 23897813). Increases cell-matrix adhesion by regulating the membrane organization of integrin alpha4/ITA4 (PubMed: 24623721 , PubMed: 8757325). Modulates adhesion and suppresses cell migration through other integrins such as the alpha6/ITGA6 and beta1/ITGB1 (PubMed: 15557282 , PubMed: 17560548). Decreases

cell-associated plasminogen activation by interfering with the interaction between urokinase-type plasminogen activator/PLAU and its receptor PLAUR (PubMed:[15677461](#)). Associates with CD4 or CD8 and delivers costimulatory signals for the TCR/CD3 pathway. Plays a role in TLR9 trafficking to acidified CpG-containing compartments by controlling interaction between TLR9 and VAMP3 and subsequent myddosome assembly (By similarity). Inhibits LPS-induced inflammatory response by preventing binding of LPS to TLR4 on the cell surface (PubMed:[36945827](#)). Plays a role in the activation of macrophages into anti-inflammatory phenotypes (By similarity). Independently of Toll- like receptor (TLR) signaling, is recruited to pathogen-containing phagosomes prior to fusion with lysosomes and thereby participates in antigen presentation (By similarity). Also acts to control angiogenesis and switch angiogenic milieu to quiescent state by binding and sequestering VEGFA and PDGFB to inhibit the signaling they trigger via their respective cell surface receptor (PubMed:[34530889](#)).


Cellular Location

Cell membrane {ECO:0000269 | PubMed:19497983, ECO:0000269 | PubMed:23897813, ECO:0000269 | PubMed:30463011, ECO:0000269 | PubMed:34530889, ECO:0000269 | PubMed:8757325, ECO:0000269 | Ref.4}; Multi-pass membrane protein Cytoplasmic vesicle, phagosome {ECO:0000250 | UniProtKB:P40237}

Tissue Location

Lymphoid specific.

Images



Western blot analysis of CD82 expression in Jurkat (A) whole cell lysates. (Predicted band size: 30 kD; Observed band size: 30-60 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.