

Recombinant Anti-CD97 alpha Rabbit mAb

Catalog # AP95105

Product Information

Application	WB, IHC, FC, IP, IF/IC
Primary Accession	P48960
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal
Calculated MW	91869


Additional Information

Gene ID	976
Other Names	CD97 antigen; Leukocyte antigen CD97; CD97
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human CD97 alpha protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	ADGRE5 (HGNC:1711)
Function	Receptor potentially involved in both adhesion and signaling processes early after leukocyte activation. Plays an essential role in leukocyte migration.
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:Q9Z0M6}; Multi-pass membrane protein
Tissue Location	Broadly expressed, found on most hematopoietic cells, including activated lymphocytes, monocytes, macrophages, dendritic cells, and granulocytes. Expressed also abundantly by smooth muscle cells. Expressed in thyroid, colorectal, gastric, esophageal and pancreatic carcinomas too. Expression are increased under inflammatory conditions in the CNS of multiple sclerosis and in synovial tissue of patients with rheumatoid arthritis. Increased expression of CD97 in the synovium is accompanied by detectable levels of soluble CD97 in the synovial fluid

Images



Western blot analysis of CD97 alpha expression in U937 (A) whole cell lysates. (Predicted band size: 92 kD; Observed band size: 70-90 kD)



Immunofluorescent analysis of CD97 alpha staining in U937 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.