

Recombinant Anti-TRPM7 Rabbit mAb

Catalog # AP95113

Product Information

Application	WB, FC, IF/IC
Primary Accession	Q96QT4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	212697

Additional Information

Gene ID	54822
Other Names	CHAK1; LTRPC7; Transient receptor potential cation channel subfamily M member 7; Channel-kinase 1; Long transient receptor potential channel 7; LTrpC-7; LTrpC7
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human TRPM7 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	TRPM7
Synonyms	CHAK1, LTRPC7 {ECO:0000303 PubMed:113855}
Function	Bifunctional protein that combines an ion channel with an intrinsic kinase domain, enabling it to modulate cellular functions either by conducting ions through the pore or by phosphorylating downstream proteins via its kinase domain. The channel is highly permeable to divalent cations, specifically calcium (Ca ²⁺), magnesium (Mg ²⁺) and zinc (Zn ²⁺) and mediates their influx (PubMed: 11385574 , PubMed: 12887921 , PubMed: 15485879 , PubMed: 24316671 , PubMed: 35561741 , PubMed: 36027648). Controls a wide range of biological processes such as Ca ²⁺ (+), Mg ²⁺ (+) and Zn ²⁺ (+) homeostasis, vesicular Zn ²⁺ (+) release channel and intracellular Ca ²⁺ (+) signaling, embryonic development, immune responses, cell motility, proliferation and differentiation (By similarity). The C-terminal alpha-kinase domain autophosphorylates cytoplasmic residues of TRPM7 (PubMed: 18365021). In vivo, TRPM7 phosphorylates SMAD2, suggesting that TRPM7 kinase may play a

role in activating SMAD signaling pathways. In vitro, TRPM7 kinase phosphorylates ANXA1 (annexin A1), myosin II isoforms and a variety of proteins with diverse cellular functions (PubMed:[15485879](#), PubMed:[18394644](#)).


Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q923J1}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Note=Localized largely in intracellular Zn(2+)-storage vesicles. {ECO:0000250|UniProtKB:Q923J1}

Images



Western blot analysis of TRPM7 expression in HeLa (A) whole cell lysates. (Predicted band size: 213 kD; Observed band size: 213 kD)



Immunofluorescent analysis of TRPM7 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.