

Recombinant Anti-Nogo Rabbit mAb

Catalog # AP95126

Product Information

Application	WB, IHC, FC, IP, IF/IC
Primary Accession	Q9NQC3
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	129931

Additional Information

Gene ID	57142
Other Names	KIAA0886; NOGO; Reticulon-4; Foocen; Neurite outgrowth inhibitor; Nogo protein; Neuroendocrine-specific protein; NSP; Neuroendocrine-specific protein C homolog; RTN-x; Reticulon-5
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human Nogo protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	RTN4 (HGNC:14085)
Function	Required to induce the formation and stabilization of endoplasmic reticulum (ER) tubules (PubMed: 24262037 , PubMed: 25612671 , PubMed: 27619977). They regulate membrane morphogenesis in the ER by promoting tubular ER production (PubMed: 24262037 , PubMed: 25612671 , PubMed: 27619977 , PubMed: 27786289). They influence nuclear envelope expansion, nuclear pore complex formation and proper localization of inner nuclear membrane proteins (PubMed: 26906412). However each isoform have specific functions mainly depending on their tissue expression specificities (Probable).
Cellular Location	[Isoform A]: Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein; Cytoplasmic side Synapse {ECO:0000250 UniProtKB:Q99P72}. Note=Anchored to the membrane of the endoplasmic reticulum (ER) through 2 putative transmembrane domains. Localizes throughout the ER tubular network (PubMed:27619977) Co-localizes with TMEM33 at the ER sheets [Isoform C]:

Endoplasmic reticulum membrane; Multi-pass membrane protein


Tissue Location

Isoform A: is specifically expressed in brain and testis and weakly in heart and skeletal muscle. Isoform B: widely expressed except for the liver. Highly expressed in endothelial cells and vascular smooth muscle cells, including blood vessels and mesenteric arteries (PubMed:15034570, PubMed:21183689). Isoform C: is expressed in brain, skeletal muscle and adipocytes. Isoform D is testis-specific.

Images



Western blot analysis of Nogo expression in HeLa (A) whole cell lysates. (Predicted band size: 130 kD; Observed band size: 40-55 kD)



Immunofluorescent analysis of Nogo staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.