

Recombinant Anti-USP7 Rabbit mAb

Catalog # AP95127

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	Q93009
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	128302

Additional Information

Gene ID	7874
Other Names	HAUSP; Ubiquitin carboxyl-terminal hydrolase 7; Deubiquitinating enzyme 7; Herpesvirus-associated ubiquitin-specific protease; Ubiquitin thioesterase 7; Ubiquitin-specific-processing protease 7
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human USP7 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	USP7 {ECO:0000303 PubMed:12093161, ECO:0000312 HGNC:HGNC:12630}
Function	Hydrolase that deubiquitinates target proteins such as ARMC5, FOXO4, DEPTOR, KAT5, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN, KMT2E/MLL5 and DAXX (PubMed:11923872, PubMed:15053880, PubMed:16964248, PubMed:18716620, PubMed:25283148, PubMed:25865756, PubMed:26678539, PubMed:28655758, PubMed:33544460, PubMed:35216969). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation (PubMed:15053880, PubMed:16845383, PubMed:18566590, PubMed:20153724). Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis (PubMed:25283148). Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis (PubMed:11923872, PubMed:26786098). Deubiquitination of

FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity (PubMed:[16964248](#)). In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML (PubMed:[18716620](#)). Deubiquitinates KMT2E/MLL5 preventing KMT2E/MLL5 proteasomal-mediated degradation (PubMed:[26678539](#)). Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UVSSA and promotes deubiquitination of ERCC6, preventing UV- induced degradation of ERCC6 (PubMed:[22466611](#), PubMed:[22466612](#)). Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1 (PubMed:[21745816](#), PubMed:[22411829](#)). Deubiquitinates alkylation repair enzyme ALKBH3. OTUD4 recruits USP7 and USP9X to stabilize ALKBH3, thereby promoting the repair of alkylated DNA lesions (PubMed:[25944111](#)). Acts as a chromatin regulator via its association with the Polycomb group (PcG) multiprotein PRC1-like complex; may act by deubiquitinating components of the PRC1-like complex (PubMed:[20601937](#)). Able to mediate deubiquitination of histone H2B; it is however unsure whether this activity takes place in vivo (PubMed:[20601937](#)). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains (PubMed:[22689415](#)). Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:[23973222](#)). Plays a role in the maintenance of the circadian clock periodicity via deubiquitination and stabilization of the CRY1 and CRY2 proteins (PubMed:[27123980](#)). Deubiquitinates REST, thereby stabilizing REST and promoting the maintenance of neural progenitor cells (PubMed:[21258371](#)). Deubiquitinates SIRT7, inhibiting SIRT7 histone deacetylase activity and regulating gluconeogenesis (PubMed:[28655758](#)). Involved in the regulation of WASH-dependent actin polymerization at the surface of endosomes and the regulation of endosomal protein recycling (PubMed:[26365382](#)). It maintains optimal WASH complex activity and precise F-actin levels via deubiquitination of TRIM27 and WASHC1 (PubMed:[26365382](#)). Mediates the deubiquitination of phosphorylated DEPTOR, promoting its stability and leading to decreased mTORC1 signaling (PubMed:[35216969](#)). Deubiquitinates H2AB1, H2BC4 and H2BC22 (By similarity).

Cellular Location


Nucleus. Cytoplasm Nucleus, PML body. Chromosome. Note=Present in a minority of ND10 nuclear bodies (PubMed:9034339). Association with ICP0/VMW110 at early times of infection leads to an increased proportion of USP7-containing ND10 (PubMed:9034339). Colocalizes with ATXN1 in the nucleus (PubMed:12093161). Colocalized with DAXX in speckled structures (PubMed:16845383). Colocalized with PML and PTEN in promyelocytic leukemia protein (PML) nuclear bodies (PubMed:18716620). Interaction with FAM170A promotes its nuclear translocation (By similarity) {ECO:0000250|UniProtKB:Q6A4J8, ECO:0000269|PubMed:12093161, ECO:0000269|PubMed:16845383, ECO:0000269|PubMed:18716620, ECO:0000269|PubMed:9034339}

Tissue Location


Expressed in neural progenitor cells (at protein level) (PubMed:21258371). Widely expressed. Overexpressed in prostate cancer.

Images

Western blot analysis of USP7 expression in Ramos (A)



whole cell lysates. (Predicted band size: 128 kD; Observed band size: 128 kD)



Immunofluorescent analysis of USP7 staining in Ramos cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.