

Recombinant Anti-BRAT1 Rabbit mAb

Catalog # AP95135

Product Information

Application	WB, FC, IP, IF/IC
Primary Accession	Q6PIG6
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	88119

Additional Information

Gene ID	221927
Other Names	BAAT1; C7orf27; BRCA1-associated ATM activator 1; BRCA1-associated protein required for ATM activation protein 1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human BRAT1 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	BRAT1 {ECO:0000303 PubMed:25657994, ECO:0000312 HGNC:HGNC:21701}
Function	Component of a multiprotein complex required for the assembly of the RNA endonuclease module of the integrator complex (PubMed: 39032489 , PubMed: 39032490). Associates with INTS9 and INTS11 in the cytoplasm and blocks the active site of INTS11 to inhibit the endonuclease activity of INTS11 before formation of the full integrator complex (PubMed: 39032489 , PubMed: 39032490). Following dissociation of WDR73 of the complex, BRAT1 facilitates the nuclear import of the INTS9-INTS11 heterodimer (PubMed: 39032489). In the nucleus, INTS4 is integrated to the INTS9-INTS11 heterodimer and BRAT1 is released from the mature RNA endonuclease module by inositol hexakisphosphate (InsP6) (PubMed: 39032489). BRAT1 is also involved in DNA damage response; activates kinases ATM, SMC1A and PRKDC by modulating their phosphorylation status following ionizing radiation (IR) stress (PubMed: 16452482 , PubMed: 22977523). Plays a role in regulating mitochondrial function and cell proliferation (PubMed: 25070371). Required for protein stability of MTOR and MTOR-related proteins, and cell

cycle progress by growth factors (PubMed:[25657994](#)).


Cellular Location

Nucleus. Cytoplasm Note=Present at double strand breaks (DSBs)following ionizing radiation treatment. The ubiquitinated form localizes in the nucleus in a NDFIP1- dependent manner.


Tissue Location

Ubiquitously expressed.

Images



Western blot analysis of BRAT1 expression in HeLa (A) whole cell lysates. (Predicted band size: 88 kD; Observed band size: 75 kD)



Immunofluorescent analysis of BRAT1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.