

Recombinant Anti-E2F6 Rabbit mAb

Catalog # AP95162

Product Information

Application	WB, FC, IF/IC
Primary Accession	O75461
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	31844

Additional Information


Gene ID	1876
Other Names	Transcription factor E2F6; E2F-6
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human E2F6 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	E2F6 {ECO:0000303 PubMed:9689056, ECO:0000312 HGNC:HGNC:3120}
Function	Inhibitor of E2F-dependent transcription (PubMed: 9501179 , PubMed: 9689056 , PubMed: 9704927). Binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' (PubMed: 9501179). Has a preference for the 5'-TTTCCCGC-3' E2F recognition site (PubMed: 9501179). E2F6 lacks the transcriptional activation and pocket protein binding domains (PubMed: 9501179 , PubMed: 9704927). Appears to regulate a subset of E2F-dependent genes whose products are required for entry into the cell cycle but not for normal cell cycle progression (PubMed: 9501179 , PubMed: 9689056). Represses expression of some meiosis-specific genes, including SLC25A31/ANT4 (By similarity). May silence expression via the recruitment of a chromatin remodeling complex containing histone H3-K9 methyltransferase activity. Overexpression delays the exit of cells from the S-phase (PubMed: 9501179).
Cellular Location	Nucleus
Tissue Location	Expressed in all tissues examined. Highest levels in placenta, skeletal muscle,

heart, ovary, kidney, small intestine and spleen.

Images



Western blot analysis of E2F6 expression in K562 (A) whole cell lysates. (Predicted band size: 32 kD; Observed band size: 32 kD)



Immunofluorescent analysis of E2F6 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.