

Recombinant Anti-DGAT1 Rabbit mAb

Catalog # AP95177

Product Information

Application	WB, FC, IP, IF/IC
Primary Accession	O75907
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	55278

Additional Information

Gene ID	8694
Other Names	AGRP1; DGAT; Diacylglycerol O-acyltransferase 1; ACAT-related gene product 1; Acyl-CoA retinol O-fatty-acyltransferase; ARAT; Retinol O-fatty-acyltransferase; Diglyceride acyltransferase
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human DGAT1 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	DGAT1 {ECO:0000303 PubMed:16214399, ECO:0000312 HGNC:HGNC:2843}
Function	Catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol and fatty acyl CoA as substrates (PubMed: 16214399 , PubMed: 18768481 , PubMed: 28420705 , PubMed: 32433610 , PubMed: 32433611 , PubMed: 9756920). Highly expressed in epithelial cells of the small intestine and its activity is essential for the absorption of dietary fats (PubMed: 18768481). In liver, plays a role in esterifying exogenous fatty acids to glycerol, and is required to synthesize fat for storage (PubMed: 16214399). Also present in female mammary glands, where it produces fat in the milk (By similarity). May be involved in VLDL (very low density lipoprotein) assembly (PubMed: 18768481). In contrast to DGAT2 it is not essential for survival (By similarity). Functions as the major acyl-CoA retinol acyltransferase (ARAT) in the skin, where it acts to maintain retinoid homeostasis and prevent retinoid toxicity leading to skin and hair disorders (PubMed: 16214399). Exhibits additional acyltransferase activities, includin acyl CoA:monoacylglycerol acyltransferase (MGAT), wax monoester and wax diester synthases (By

similarity). Also able to use 1-monoalkylglycerol (1-MAkG) as an acyl acceptor for the synthesis of monoalkyl-monoacylglycerol (MAMAG) (PubMed:[28420705](#)).


Cellular Location

Endoplasmic reticulum membrane {ECO:0000250 | UniProtKB:Q9Z2A7};
Multi-pass membrane protein

Images



Western blot analysis of DGAT1 expression in HeLa (A) whole cell lysates. (Predicted band size: 55 kD; Observed band size: 48 kD)



Immunofluorescent analysis of DGAT1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.