

Recombinant Anti-Nesprin 1 Rabbit mAb

Catalog # AP95183

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	Q8NF91
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	1011086

Additional Information

Gene ID	23345
Other Names	C6orf98; KIAA0796; KIAA1262; KIAA1756; MYNE1; Nesprin-1; Enaptin; Myocyte nuclear envelope protein 1; Myne-1; Nuclear envelope spectrin repeat protein 1; Synaptic nuclear envelope protein 1; Syne-1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human Nesprin 1 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SYNE1 (HGNC:17089)
Function	Multi-isomeric modular protein which forms a linking network between organelles and the actin cytoskeleton to maintain the subcellular spatial organization. As a component of the LINC (Linker of Nucleoskeleton and Cytoskeleton) complex involved in the connection between the nuclear lamina and the cytoskeleton. The nucleocytoplasmic interactions established by the LINC complex play an important role in the transmission of mechanical forces across the nuclear envelope and in nuclear movement and positioning. May be involved in nucleus- centrosome attachment and nuclear migration in neural progenitors implicating LINC complex association with SUN1/2 and probably association with cytoplasmic dynein-dynactin motor complexes; SYNE1 and SYNE2 may act redundantly. Required for centrosome migration to the apical cell surface during early ciliogenesis. May be involved in nuclear remodeling during sperm head formation in spermatogenesis; a probable SUN3:SYNE1/KASH1 LINC complex may tether spermatid nuclei to posterior cytoskeletal structures such as the manchette.


Cellular Location

Nucleus outer membrane; Single-pass type IV membrane protein; Cytoplasmic side Nucleus. Nucleus envelope. Cytoplasm, cytoskeleton. Cytoplasm, myofibril, sarcomere. Note=The largest part of the protein is cytoplasmic, while its C-terminal part is associated with the nuclear envelope, most probably the outer nuclear membrane. In skeletal and smooth muscles, a significant amount is found in the sarcomeres. In myoblasts, relocated from the nuclear envelope to the nucleus and cytoplasm during cell differentiation


Tissue Location

Expressed in HeLa, A431, A172 and HaCaT cells (at protein level). Widely expressed. Highly expressed in skeletal and smooth muscles, heart, spleen, peripheral blood leukocytes, pancreas, cerebellum, stomach, kidney and placenta. Isoform GSRP-56 is predominantly expressed in heart and skeletal muscle (at protein level).

Images



Western blot analysis of Nesprin 1 expression in HeLa (A) whole cell lysates. (Predicted band size: 1011 kD; Observed band size: 112 kD)



Immunofluorescent analysis of Nesprin 1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.