

Recombinant Anti-CHD1L Rabbit mAb

Catalog # AP95186

Product Information

Application	WB, IHC, FC, IP, IF/IC
Primary Accession	Q86WJ1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal
Calculated MW	101000

Additional Information

Gene ID	9557
Other Names	ALC1; Chromodomain-helicase-DNA-binding protein 1-like; Amplified in liver cancer protein 1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human CHD1L protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CHD1L {ECO:0000303 PubMed:34210977, ECO:0000312 HGNC:HGNC:1916}
Function	ATP-dependent chromatin remodeler that mediates chromatin- remodeling following DNA damage (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 33357431 , PubMed: 34210977 , PubMed: 34486521 , PubMed: 34874266). Recruited to DNA damage sites through interaction with poly-ADP-ribose: specifically recognizes and binds histones that are poly-ADP-ribosylated on serine residues in response to DNA damage (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 34486521 , PubMed: 34874266). Poly-ADP-ribose-binding activates the ATP-dependent chromatin remodeler activity, thereby regulating chromatin during DNA repair (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 34486521 , PubMed: 34874266). Catalyzes nucleosome sliding away from DNA breaks in an ATP-dependent manner (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653). Chromatin remodeling activity promotes PARP2 removal from chromatin (PubMed: 33275888).


Cellular Location

Nucleus. Chromosome Note=Localizes at sites of DNA damage; recruited by histones H2B and H3 poly-ADP-ribosylated on 'Ser-6' and 'Ser-10', respectively (H2BS6ADPr and H3S10ADPr) by PARP1 or PARP2.


Tissue Location

Frequently overexpressed in hepatomacellular carcinomas.

Images



Western blot analysis of CHD1L expression in A549 (A) whole cell lysates. (Predicted band size: 101 kD; Observed band size: 110 kD)



Immunofluorescent analysis of CHD1L staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.