

Recombinant Anti-MTH1 Rabbit mAb

Catalog # AP95197

Product Information

Application	WB, IHC, FC, IP, IF/IC
Primary Accession	P36639
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	17952

Additional Information

Gene ID	4521
Other Names	MTH1; 7, 8-dihydro-8-oxoguanine triphosphatase; 2-hydroxy-dATP diphosphatase; 8-oxo-dGTPase; Nucleoside diphosphate-linked moiety X motif 1; Nudix motif 1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human MTH1 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	NUDT1
Synonyms	MTH1 {ECO:0000303 PubMed:7713500}
Function	Oxidized purine nucleoside triphosphate hydrolase which is a prominent sanitizer of the oxidized nucleotide pool (PubMed:10608900, PubMed:12857738, PubMed:22556419, PubMed:24695224, PubMed:24695225, PubMed:26238318, PubMed:28679043, PubMed:7713500, PubMed:8226881). Catalyzes the hydrolysis of 2-oxo-dATP (2-hydroxy-dATP) into 2-oxo-dAMP (PubMed:10373420). Also has a significant hydrolase activity toward 2- oxo-ATP, 8-oxo-dGTP and 8-oxo-dATP (PubMed:10373420, PubMed:11139615). Through the hydrolysis of oxidized purine nucleoside triphosphates, prevents their incorporation into DNA and the subsequent transversions A:T to C:G and G:C to T:A (PubMed:10373420, PubMed:10608900, PubMed:11756418, PubMed:12857738, PubMed:16607562, PubMed:24695224, PubMed:24695225, PubMed:26999531, PubMed:28035004, PubMed:8226881). Also catalyzes the

hydrolysis of methylated purine nucleoside triphosphate preventing their integration into DNA (PubMed:[30304478](#), PubMed:[32144205](#)). Through this antimutagenic activity protects cells from oxidative stress (PubMed:[10608900](#), PubMed:[12857738](#), PubMed:[24695224](#), PubMed:[24695225](#), PubMed:[30304478](#), PubMed:[32144205](#), PubMed:[7713500](#), PubMed:[8226881](#)).


Cellular Location

[Isoform p18]: Cytoplasm, cytosol. Mitochondrion matrix. Nucleus.
Note=Mostly present in cytosol (PubMed:7782328). A minor proportion is mitochondrial (PubMed:7782328) A very small amount of the protein is associated with nuclei (PubMed:7782328).


Tissue Location

Widely expressed with highest expression in thymus, testis, embryo and proliferating blood lymphocytes

Images



Western blot analysis of MTH1 expression in Jurkat (A) whole cell lysates. (Predicted band size: 18 kD; Observed band size: 18 kD)



Immunofluorescent analysis of MTH1 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.