

Recombinant Anti-DHEA-ST Rabbit mAb

Catalog # AP95199

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	Q06520
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	33780

Additional Information

Gene ID	6822
Other Names	HST; STD; Bile salt sulfotransferase; Dehydroepiandrosterone sulfotransferase; DHEA-ST; Hydroxysteroid Sulfotransferase; HST; ST2; ST2A3; Sulfotransferase 2A1; ST2A1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human DHEA-ST protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SULT2A1
Synonyms	HST, STD
Function	Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the sulfonation of steroids and bile acids in the liver and adrenal glands (PubMed: 14573603 , PubMed: 18042734 , PubMed: 19589875 , PubMed: 20102295 , PubMed: 21187059 , PubMed: 2268288 , PubMed: 29671343 , PubMed: 7678732 , PubMed: 7854148). Mediates the sulfation of a wide range of steroids and sterols, including pregnenolone, androsterone, DHEA, bile acids, cholesterol and as well many xenobiotics that contain alcohol and phenol functional groups (PubMed: 14573603 , PubMed: 18042734 , PubMed: 19589875 , PubMed: 20102295 , PubMed: 21187059 , PubMed: 2268288 , PubMed: 29671343 , PubMed: 7678732 , PubMed: 7854148). Sulfonation increases the water solubility of most compounds, and therefore their renal excretion, but it can also result in bioactivation to form active metabolites. Plays an important role in

maintaining steroid and lipid homeostasis (PubMed:[14573603](#), PubMed:[19589875](#), PubMed:[21187059](#)). Plays a key role in bile acid metabolism, mediating formation of 3-sulfated bile acids (PubMed:[2268288](#), PubMed:[20102295](#)). In addition, catalyzes the metabolic activation of potent carcinogenic polycyclic arylmethanols (By similarity).

Cellular Location

Cytoplasm.


Tissue Location

Liver, adrenal and at lower level in the kidney. Is present in human fetus in higher level in the adrenal than the liver and the kidney

Images



Western blot analysis of DHEA-ST expression in HepG2 (A) whole cell lysates. (Predicted band size: 34 kD; Observed band size: 34 kD)



Immunofluorescent analysis of DHEA-ST staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.