

Recombinant Anti-BAZ1B Rabbit mAb

Catalog # AP95204

Product Information

Application	WB, FC
Primary Accession	Q9UIG0
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	170903

Additional Information

Gene ID	9031
Other Names	WBSC10; WBSCR10; WBSCR9; WSTF; Tyrosine-protein kinase BAZ1B; Bromodomain adjacent to zinc finger domain protein 1B; Williams syndrome transcription factor; Williams-Beuren syndrome chromosomal region 10 protein; Williams-Beuren syndrome chromosomal region 9 protein; hWALp2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human BAZ1B protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	BAZ1B
Synonyms	WBSC10, WBSCR10, WBSCR9, WSTF
Function	Atypical tyrosine-protein kinase that plays a central role in chromatin remodeling and acts as a transcription regulator (PubMed: 19092802). Involved in DNA damage response by phosphorylating 'Tyr-142' of histone H2AX (H2AXY142ph) (PubMed: 19092802 , PubMed: 19234442). H2AXY142ph plays a central role in DNA repair and acts as a mark that distinguishes between apoptotic and repair responses to genotoxic stress (PubMed: 19092802 , PubMed: 19234442). Regulatory subunit of the ATP-dependent WICH-1 and WICH-5 ISWI chromatin remodeling complexes, which form ordered nucleosome arrays on chromatin and facilitate access to DNA during DNA-templated processes such as DNA replication, transcription, and repair (PubMed: 11980720 , PubMed: 28801535). Both complexes regulate the spacing of nucleosomes along the chromatin and have the ability to slide

mononucleosomes to the center of a DNA template (PubMed:[28801535](#)). The WICH-1 ISWI chromatin remodeling complex has a lower ATP hydrolysis rate than the WICH-5 ISWI chromatin remodeling complex (PubMed:[28801535](#)). The WICH-5 ISWI chromatin-remodeling complex regulates the transcription of various genes, has a role in RNA polymerase I transcription (By similarity). Within the B-WICH complex has a role in RNA polymerase III transcription (PubMed:[16603771](#)). Mediates the recruitment of the WICH-5 ISWI chromatin remodeling complex to replication foci during DNA replication (PubMed:[15543136](#)).


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00063, ECO:0000255 | PROSITE-ProRule:PRU00475, ECO:0000269 | PubMed:11980720, ECO:0000269 | PubMed:15543136, ECO:0000269 | PubMed:16603771, ECO:0000269 | PubMed:25593309}. Note=Accumulates in pericentromeric heterochromatin during replication (PubMed:15543136). Co-localizes with PCNA at replication foci during S phase (PubMed:15543136). Co-localizes with SMARCA5/SNF2H at replication foci during late-S phase (PubMed:15543136). Also localizes to replication foci independently of SMARCA5/SNF2H and PCNA (PubMed:15543136). Localizes to sites of DNA damage (PubMed:25593309).

Tissue Location

Ubiquitously expressed with high levels of expression in heart, brain, placenta, skeletal muscle and ovary

Images



Western blot analysis of BAZ1B expression in SHSY5Y (A) whole cell lysates. (Predicted band size: 171 kD; Observed band size: 185 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.