

Recombinant Anti-RBP2 Rabbit mAb

Catalog # AP95208

Product Information

Application	WB, FC, IP, IF/IC
Primary Accession	P29375
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal
Calculated MW	192095

Additional Information

Gene ID	5927
Other Names	JARID1A; RBBP2; RBP2; Lysine-specific demethylase 5A; Histone demethylase JARID1A; Jumonji/ARID domain-containing protein 1A; Retinoblastoma-binding protein 2; RBBP-2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human RBP2 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	KDM5A (HGNC:9886)
Function	Histone demethylase that specifically demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9', H3 'Lys-27', H3 'Lys-36', H3 'Lys-79' or H4 'Lys-20'. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-4'. Regulates specific gene transcription through DNA-binding on 5'-CCGCCC-3' motif (PubMed:18270511). May stimulate transcription mediated by nuclear receptors. Involved in transcriptional regulation of Hox proteins during cell differentiation (PubMed:19430464). May participate in transcriptional repression of cytokines such as CXCL12. Plays a role in the regulation of the circadian rhythm and in maintaining the normal periodicity of the circadian clock. In a histone demethylase-independent manner, acts as a coactivator of the CLOCK-BMAL1-mediated transcriptional activation of PER1/2 and other clock-controlled genes and increases histone acetylation at PER1/2 promoters by inhibiting the activity of HDAC1 (By similarity). Seems to act as a transcriptional corepressor for some genes such as MT1F and to favor the

proliferation of cancer cells (PubMed:[27427228](#)).


Cellular Location

Nucleus, nucleolus. Nucleus {ECO:0000250|UniProtKB:Q3UXZ9}
Note=Occupies promoters of genes involved in RNA metabolism and mitochondrial function. {ECO:0000250|UniProtKB:Q3UXZ9}

Images



Western blot analysis of RBP2 expression in HEK293 (A) whole cell lysates. (Predicted band size: 192 kD; Observed band size: 192 kD)



Immunofluorescent analysis of RBP2 staining in HEK293 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.