

Recombinant Anti-WASP Rabbit mAb

Catalog # AP95212

Product Information

Application	WB, FC, IP, IF/IC
Primary Accession	P42768
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	52913

Additional Information

Gene ID	7454
Other Names	IMD2; Wiskott-Aldrich syndrome protein; WASp
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human WASP protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	WAS
Synonyms	IMD2
Function	Effector protein for Rho-type GTPases that regulates actin filament reorganization via its interaction with the Arp2/3 complex (PubMed: 12235133 , PubMed: 12769847 , PubMed: 16275905). Important for efficient actin polymerization (PubMed: 12235133 , PubMed: 16275905 , PubMed: 8625410). Possible regulator of lymphocyte and platelet function (PubMed: 9405671). Mediates actin filament reorganization and the formation of actin pedestals upon infection by pathogenic bacteria (PubMed: 18650809). In addition to its role in the cytoplasmic cytoskeleton, also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed: 20574068). Promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed: 29925947).
Cellular Location	Cytoplasm, cytoskeleton. Nucleus


Tissue Location

Expressed predominantly in the thymus. Also found, to a much lesser extent, in the spleen.

Images



Western blot analysis of WASP expression in Jurkat (A) whole cell lysates. (Predicted band size: 53 kD; Observed band size: 53 kD)



Immunofluorescent analysis of WASP staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.