10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Recombinant Anti-Synoviolin Rabbit mAb

Catalog # AP95320

Product Information

Application	WB, FC, IP, IF/IC
Primary Accession	Q86TM6
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal
Calculated MW	67685

Additional Information

Gene ID	84447
Other Names	HRD1; KIAA1810; E3 ubiquitin-protein ligase synoviolin; Synovial apoptosis inhibitor 1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human Synoviolin protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SYVN1 {ECO:0000303|PubMed:15489334}
Function	E3 ubiquitin-protein ligase which accepts ubiquitin specifically from endoplasmic reticulum-associated UBC7 E2 ligase and transfers it to substrates, promoting their degradation (PubMed:12459480, PubMed:12646171, PubMed:12975321, PubMed:14593114, PubMed:16289116, PubMed:16847254, PubMed:17059562, PubMed:17141218, PubMed:17170702, PubMed:22607976, PubMed:27827840, PubMed:26471130, PubMed:28827405). Component of the endoplasmic reticulum quality control (ERQC) system also called ER-associated degradation (ERAD) involved in ubiquitin-dependent degradation of misfolded endoplasmic reticulum proteins (PubMed:12459480, PubMed:12646171, PubMed:12975321, PubMed:14593114, PubMed:16289116, PubMed:16847254, PubMed:17059562, PubMed:17141218, PubMed:17170702, PubMed:22607976, PubMed:26471130, PubMed:28842558). Also promotes the degradation of normal but naturally short-lived proteins such as SGK. Protects cells from ER stress-induced apoptosis. Protects neurons from

apoptosis induced by polyglutamine- expanded huntingtin (HTT) or unfolded GPR37 by promoting their degradation (PubMed:17141218). Sequesters p53/TP53 in the cytoplasm and promotes its degradation, thereby negatively regulating its biological function in transcription, cell cycle regulation and apoptosis (PubMed:17170702). Mediates the ubiquitination and subsequent degradation of cytoplasmic NFE2L1 (By similarity). During the early stage of B cell development, required for degradation of the pre-B cell receptor (pre-BCR) complex, hence supporting further differentiation into mature B cells (By similarity).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein


Tissue Location

Ubiquitously expressed, with highest levels in liver and kidney (at protein level). Up-regulated in synovial tissues from patients with rheumatoid arthritis (at protein level)

Images



Western blot analysis of Synoviolin expression in Hela (A) whole cell lysates. (Predicted band size: 68 kD; Observed band size: 65 kD)



Immunofluorescent analysis of Synoviolin staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.