

Recombinant Anti-DFF45 Rabbit mAb

Catalog # AP95331

Product Information

Application	WB, IHC, FC, IP, IF/IC
Primary Accession	O00273
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	36522

Additional Information



Gene ID	1676
Other Names	DFF1; DFF45; DNA fragmentation factor subunit alpha; DNA fragmentation factor 45 kDa subunit; DFF-45; Inhibitor of CAD; ICAD
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human DFF45 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	DFFA
Synonyms	DFF1, DFF45
Function	Inhibitor of the caspase-activated DNase (DFF40).
Cellular Location	Cytoplasm.

Images

Western blot analysis of DFF45 expression in HeLa (A) whole cell lysates. (Predicted band size: 37 kD; Observed band size: 37 kD)



Immunofluorescent analysis of DFF45 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.