

Recombinant Anti-UBE4A Rabbit mAb

Catalog # AP95366

Product Information

Application	WB, FC, IF/IC
Primary Accession	Q14139
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	122561

Additional Information


Gene ID	9354
Other Names	KIAA0126; Ubiquitin conjugation factor E4 A
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within human UBE4A protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IF/IC~~N/A
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	UBE4A (HGNC:12499)
Function	Ubiquitin-protein ligase that probably functions as an E3 ligase in conjunction with specific E1 and E2 ligases. May also function as an E4 ligase mediating the assembly of polyubiquitin chains on substrates ubiquitinated by another E3 ubiquitin ligase. Mediates 'Lys-48'-linked polyubiquitination of substrates.
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:E9Q735}.

Images

Western blot analysis of UBE4A expression in HEK293 (A) whole cell lysates. (Predicted band size: 123 kD; Observed band size: 123 kD)



Immunofluorescent analysis of UBE4A staining in HEK293 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.