

Recombinant Anti-c-CBL (Phospho-Y774) Rabbit mAb

Catalog # AP95378

Product Information

Application	WB, IHC, FC
Primary Accession	P22681
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	99633

Additional Information

Gene ID	867
Other Names	CBL2; RNF55; E3 ubiquitin-protein ligase CBL; Casitas B-lineage lymphoma proto-oncogene; Proto-oncogene c-Cbl; RING finger protein 55; Signal transduction protein CBL
Target/Specificity	KLH-conjugated synthetic Phospho-peptide corresponding to residues surrounding Y774 of human c-CBL. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~1:10~50
Format	Liquid in PBS, pH 7.3, 50% glycerol, 0.05% BSA, and 0.05% Proclin300.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	CBL
Synonyms	CBL2, RNF55
Function	E3 ubiquitin-protein ligase that acts as a negative regulator of many signaling pathways by mediating ubiquitination of cell surface receptors (PubMed:10514377, PubMed:11896602, PubMed:14661060, PubMed:14739300, PubMed:15190072, PubMed:17509076, PubMed:18374639, PubMed:19689429, PubMed:21596750, PubMed:28381567, PubMed:40101708). Accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome (PubMed:10514377, PubMed:14661060, PubMed:14739300, PubMed:17094949, PubMed:17509076, PubMed:17974561). Recognizes activated receptor tyrosine kinases, including KIT, FLT1, FGFR1, FGFR2, PDGFRA, PDGFRB, CSF1R, EPHA8 and KDR and mediates their ubiquitination to terminate signaling (PubMed:15190072, PubMed:18374639, PubMed:21596750). Recognizes membrane-bound HCK,

SRC and other kinases of the SRC family and mediates their ubiquitination and degradation (PubMed:[11896602](#)). Ubiquitinates EGFR and SPRY2 (PubMed:[17094949](#), PubMed:[17974561](#)). Involved in LAG3-mediated inhibition of TCR signaling; following ligand-binding to LAG3, catalyzes 'Lys-63'-linked ubiquitination of LAG3, unleashing the LAG3 C-terminus from the membrane, and initiating a signaling that prevents TCR activation (PubMed:[40101708](#)). Ubiquitinates NECTIN1 following association between NECTIN1 and herpes simplex virus 1/HHV-1 envelope glycoprotein D, leading to NECTIN1 removal from cell surface (PubMed:[28381567](#)). Participates in signal transduction in hematopoietic cells. Plays an important role in the regulation of osteoblast differentiation and apoptosis (PubMed:[15190072](#), PubMed:[18374639](#)). Essential for osteoclastic bone resorption (PubMed:[14739300](#)). The 'Tyr-731' phosphorylated form induces the activation and recruitment of phosphatidylinositol 3-kinase to the cell membrane in a signaling pathway that is critical for osteoclast function (PubMed:[14739300](#)). In association with CBLB, required for proper feedback inhibition of ciliary platelet-derived growth factor receptor-alpha (PDGFRA) signaling pathway via ubiquitination and internalization of PDGFRA (By similarity).

Cellular Location

Cytoplasm. Cell membrane. Cell projection, cilium. Golgi apparatus.
Note=Colocalizes with FGFR2 in lipid rafts at the cell membrane

Images



Western blot analysis of c-CBL (Phospho-Y774) expression in Jurkat treated with pervanadate (A) whole cell lysates. (Predicted band size: 100 kD; Observed band size: 120 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.