

Anti-CD13 Antibody

Catalog # AP95476

Product Information

Application	WB, FC, IF/IC
Primary Accession	P15144
Reactivity	Human
Host	Mouse
Clonality	Monoclonal (clone: 22A5)
Calculated MW	109540

Additional Information


Gene ID	290
Other Names	APN; CD13; PEPN; Aminopeptidase N; AP-N; hAPN; Alanyl aminopeptidase; Aminopeptidase M; AP-M; Microsomal aminopeptidase; Myeloid plasma membrane glycoprotein CD13; gp150; CD antigen CD13
Target/Specificity	Native purified human CD13.
Dilution	WB~~WB (1/500 - 1/1000) FC~~FC (1/500 - 1/2000) IF/IC~~N/A
Format	Mouse IgG2a. Liquid in PBS, pH 7.3, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	ANPEP
Synonyms	APN, CD13, PEPN
Function	Broad specificity aminopeptidase which plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Also involved in the processing of various peptides including peptide hormones, such as angiotensin III and IV, neuropeptides, and chemokines. May also be involved the cleavage of peptides bound to major histocompatibility complex class II molecules of antigen presenting cells. May have a role in angiogenesis and promote cholesterol crystallization. May have a role in amino acid transport by acting as binding partner of amino acid transporter SLC6A19 and regulating its activity (By similarity).
Cellular Location	Cell membrane; Single-pass type II membrane protein. Note=Also found as a soluble form
Tissue Location	Expressed in epithelial cells of the kidney, intestine, and respiratory tract;

granulocytes, monocytes, fibroblasts, endothelial cells, cerebral pericytes at the blood-brain barrier, synaptic membranes of cells in the CNS. Also expressed in endometrial stromal cells, but not in the endometrial glandular cells. Found in the vasculature of tissues that undergo angiogenesis and in malignant gliomas and lymph node metastases from multiple tumor types but not in blood vessels of normal tissues. A soluble form has been found in plasma. It is found to be elevated in plasma and effusions of cancer patients.

Images



Western blot analysis of CD13 expression in A2780 (A), U87 (B), A375 (C), mouse liver (D) whole cell lysates. (Predicted band size: 109 kD; Observed band size: 160 kD)



Flow cytometric analysis of human peripheral blood granulocytes using Anti-CD13 Antibody, followed by anti-mouse IgG PE.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.